arXiv:1102.2509v1 [math.FA] 12 Feb 2011

Approximation properties for Baskakov-Kantorovich-Stancu type operators based on $q-$ integers

Çiğdem Atakut[1], İbrahim Büyükyazıcı[2]

[1]Ankara University, Faculty of Science, Department of Mathematics, Tandogan
06100, Ankara, Turkey.
e-mail: atakut@science.ankara.edu.tr

[2]Gazi University, Department of Mathematics, Çubuk, Ankara, Turkey.
e-mail: bibrahim@gazi.edu.tr

Abstract: In this paper, we give an interesting generalization of the Stancu type Baskakov-Kantorovich operators based on the $q-$integers and investigate their approximation properties. Also, we obtain the estimates for the rate of convergence for a sequence of them by the weighted modulus of smoothness.

Key words: $q-$integer; $q-$ Baskakov-Kantorovich operators; Baskakov-Kantorovich-Stancu operators; Weighted spaces; Rate of convergence; Weighted modulus of smoothness.

2000 MSC: 41A10; 41A25; 41A36.

1 Introduction

In recent years, due to the intensive development of $q-$ calculus, generalizations of some operators related to $q-$ calculus have emerged (see $[2,5,6,10-18]$). Aral and Gupta defined $q-$ generalization of the Baskakov operator and investigated approximation properties of these operators in [2]. In [12], Gupta and Radu introduced the Baskakov- Kantorovich operators based on $q-$integers and investigated their weighted statistical approximation properties. They also proved some direct estimations for error using weighted modulus of smoothness in case $0<q<1$. In recent study Büyükyazıcı and Atakut [5] introduced a new Stancu type generalization of $q-$ Baskakov operator is defined as

$$L_n^{\alpha,\beta}(f;q,x)=\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \quad f(\frac{1}{q^{k-1}}\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}) \tag{1}$$

where $0\leq\alpha\leq\beta$, $q\in(0,1)$, $f\in C[0,\infty)$ and the following conditions are provided:

Let $\{\varphi_n\}$ $(n=1,2,...)$ $\varphi_n:\mathbb{R}\to\mathbb{R}$ be a sequence which is satisfying following conditions,

(i) φ_n $(n = 1, 2, ...)$, $k-$ times continuously $q-$ differentiable any closed interval $[0, b]$,

(ii) $\varphi_n(0) = 1, (n = 1, 2, ...)$,

(iii) for all $x \in [0, b]$, and $(k = 0, 1, ...; n = 1, 2, ...)$, $(-1)^k D_q^k (\varphi_n(x)) \geq 0$,

(iv) there exists a positive integer $m(n)$, such that

$$D_q^k (\varphi_n(x)) = - [n]_q D_q^{k-1} \varphi_{m(n)}(x)(1 + \alpha_{k,n}(x)), \tag{2}$$

$(k = 0, 1, ...; n = 1, 2, ...)$ and $x \in [0, b]$ where $\alpha_{k,n}(x)$ converges to zero for $n \to \infty$ uniformly in k.

(v) $\lim\limits_{n\to\infty} \frac{[n]_q}{[m(n)]_q} = 1$.

Now, to explain the construction of the new $q-$ operators, we mention some basic definitions of $q-$ calculus and Lemma.

Let $q > 0$. For each nonnegative integer n, we define the $q-$ integer $[n]_q$ as

$$[n]_q = \begin{cases} (1 - q^n)/(1 - q) & \text{if} \quad q \neq 1 \\ n & \text{if} \quad q = 1 \end{cases}$$

and the $q-$ factorial $[n]_q!$ as

$$[n]_q! = \begin{cases} [n]_q [n-1]_q \cdots [1]_q & \text{if} \quad n \geq 1 \\ 1 & \text{if} \quad n = 0 \end{cases}$$

For the integers n and k, with $0 \leq k \leq n$, the $q-$ binomial coefficients are then defined as follows (see [14]):

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}.$$

Note that the following relation is satisfied

$$[n]_q = [n-1]_q + q^{n-1}.$$

Definition 1 *The $q-$ derivative of a function f with respect to x is*

$$D_q (f(x)) = \frac{f(qx) - f(x)}{qx - x}$$

which is also known as the Jackson derivative. High $q-$ derivatives are

$$D_q^0 (f(x)) = f(x) \ , \ D_q^n (f(x)) = D_q \left(D_q^{n-1} (f(x)) \right) \ , \ n = 1, 2, 3, ...$$

Note that as $q \to 1$, the $q-$ derivative approach the usual derivative.

Definition 2 *The $q-$integration is defined as*

$$\int_0^a f(t) d_q t = (1 - q) a \sum_{j=0}^{\infty} f(q^j a) q^j \qquad , \quad a > 0.$$

Over a general interval $[a,b]$, $0<a<b$, *one defines*

$$\int_a^b f(t)d_qt=\int_0^b f(t)d_qt-\int_0^a f(t)d_qt.$$

Definition 3 *Let* $f(x)$ *be a continuous function on some interval* $[a,b]$ *and* $c\in(a,b)$. *Jackson's* $q-$ *Taylor formula (see* [13, 14]*) is given by*

$$f(x)=\sum_{k=0}^{\infty}\frac{(D_q^kf)(c)}{[k]_q!}(x-c)_q^k$$

where $(x-c)_q^k=\prod_{i=0}^{k-1}(x-cq^i)$.

First we need the following auxiliary result. Throughout the paper, we use e_i the test functions defined by $e_i(t):=t^i$ for every integer $i\geq 0$.

Lemma 4 (from [5]) *For* $L_n^{\alpha,\beta}(e_i(t);q,x)$, $i=0,1,2$ *the following identities hold:*

$$L_n^{\alpha,\beta}(e_0;q,x)=1, \tag{3}$$

$$L_n^{\alpha,\beta}(e_1;q,x)=\frac{[n]_q}{[n]_q+\beta}x(1+\alpha_{1,n}(x))+\frac{\alpha}{[n]_q+\beta}, \tag{4}$$

$$\begin{aligned} L_n^{\alpha,\beta}(e_2;q,x) &= \frac{[n]_q\,[m(n)]_q}{q\left([n]_q+\beta\right)^2}x^2(1+\alpha_{1,m(n)}(x))(1+\alpha_{2,n}(x)) \\ &\quad +\frac{[n]_q\,(2\alpha+1)}{\left([n]_q+\beta\right)^2}x\,(1+\alpha_{1,n}(x))+\frac{\alpha^2}{\left([n]_q+\beta\right)^2}. \end{aligned} \tag{5}$$

2 Some properties of Stancu type q-Baskakov-Kantorovich operators

In addition to the above conditions $(i)-(v)$, $\varphi_n(x)$ and $\alpha_{k,n}(x)$ are satisfied following condition:

$$(vi)\quad \varphi_n(x)(1+\alpha_{0,n}(x))\leq 1\ ,\quad \text{for all } x\in[0,b]\,,(n=1,2,...).$$

In this paper, under the conditions $(i)-(vi)$, we definition a new generalization of Stancu type $q-$Baskakov-Kantorovich operators as following

$$L_n^{*(\alpha,\beta)}(f;q,x)=\left([n]_q+\beta\right)\sum_{k=0}^{\infty}q^{\frac{k(k-1)}{2}}\frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k\int\limits_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}}f\left(q^{-k+1}t\right)d_qt\quad, \tag{6}$$

where $x\in\mathbb{R}_+$, $n\in N$, $0\leq\alpha\leq\beta$.

Note that, when $q=1$, the operators given by (6) is reduced to the Kantorovich-Baskakov-Stancu type operators (see [3]) and if we choose $q=1$, $\varphi_n(x)=(1+x)^{-n}$ and $\alpha=\beta=0$, we obtain Baskakov-Kantorovich operators (see [1]).

In each of the following theorems, we assume that $q=q_n$, where $\{q_n\}$ is a sequence of real numbers such that $0<q_n<1$ for all n and $\lim_{n\to\infty}q_n=1$.

Now we give the following Lemmas, which are necessary to prove our theorems:

Lemma 5 *The following relations are satisfied:*

$$\int\limits_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}}d_qt=\frac{1}{[n]_q+\beta}, \tag{7}$$

$$\int\limits_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}}td_qt=\frac{[2]_q[k]_q+q^k(1+2\alpha)}{[2]_q\left([n]_q+\beta\right)^2}\ , \tag{8}$$

$$\int\limits_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}}t^2d_qt=\frac{[3]_q[k]_q^2+q^k[k]_q\left((1+3\alpha)[2]_q+1\right)+\left(1+3\alpha+3\alpha^2\right)q^{2k}}{[3]_q\left([n]_q+\beta\right)^3}. \tag{9}$$

Proof. From properties of $q-$analogue integration, by simple computation we obtain $(7-9)$. ■

By the following Lemma Korovkin's conditions are satisfied.

Lemma 6 *For all* $x\in\mathbb{R}_+$, $n\in N$, $\alpha,\beta\geq0$ *and* $0<q<1$, *we have*

$$L_n^{*(\alpha,\beta)}(e_0;q,x)=1, \tag{10}$$

$$L_n^{*(\alpha,\beta)}(e_1;q,x)=\frac{[n]_q}{[n]_q+\beta}x\left(1+\alpha_{1,n}(x)\right)+\frac{q(1+2\alpha)}{[2]_q\left([n]_q+\beta\right)}\ , \tag{11}$$

$$\begin{aligned} L_n^{*(\alpha,\beta)}(e_2;q,x) &= \frac{[n]_q\,[m(n)]_q}{q\left([n]_q+\beta\right)^2}\left(1+\alpha_{1,m(n)}(x)\right)\left(1+\alpha_{2,n}(x)\right)x^2 \\ &\quad +\frac{[n]_q\left[[3]_q+q\left((1+3\alpha)\,[2]_q+1\right)\right]}{[3]_q\left([n]_q+\beta\right)^2}\left(1+\alpha_{1,n}(x)\right)x+\frac{q^2\left(1+3\alpha+3\alpha^2\right)}{[3]_q\left([n]_q+\beta\right)^2}. \end{aligned} \tag{12}$$

Proof. From definition (6) and the identities (3) and (7), we can easily obtain

$$\begin{aligned} L_n^{*(\alpha,\beta)}(e_0;q,x) &= \left([n]_q+\beta\right)\sum_{k=0}^{\infty}q^{\frac{k(k-1)}{2}}\frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k\int_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}}d_qt \\ &= L_n^{\alpha,\beta}(e_0;q,x)=1. \end{aligned}$$

Now for e_1,from (3), (4) and (8) we can write

$$\begin{aligned} L_n^{*(\alpha,\beta)}(e_1;q,x) &= \left([n]_q+\beta\right)\sum_{k=0}^{\infty}q^{\frac{k(k-1)}{2}}\frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k\int_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}}q^{-k+1}t\,d_qt \\ &= \sum_{k=0}^{\infty}q^{\frac{k(k-1)}{2}}\frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}\sum_{k=0}^{\infty}q^{\frac{k(k-1)}{2}}\frac{[k]_q+q^{k-1}\alpha}{q^{k-1}\left([n]_q+\beta\right)} \\ &\quad -\sum_{k=0}^{\infty}q^{\frac{k(k-1)}{2}}\frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}\frac{q^{k-1}\alpha}{q^{k-1}\left([n]_q+\beta\right)} \\ &\quad +\frac{q(1+2\alpha)}{[2]_q\left([n]_q+\beta\right)}\sum_{k=0}^{\infty}q^{\frac{k(k-1)}{2}}\frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \\ &= L_n^{\alpha,\beta}(e_1;q,x)-\frac{\alpha}{[n]_q+\beta}L_n^{\alpha,\beta}(e_0;q,x)+\frac{q(1+2\alpha)}{[2]_q\left([n]_q+\beta\right)}L_n^{\alpha,\beta}(e_0;q,x) \\ &= \frac{[n]_q}{[n]_q+\beta}x\left(1+\alpha_{1,n}(x)\right)+\frac{q(1+2\alpha)}{[2]_q\left([n]_q+\beta\right)}. \end{aligned}$$

The finally, for e_2, we use (3), (4), (5) and (9), one has

$$L_n^{*(\alpha,\beta)}(e_2;q,x) = \left([n]_q+\beta\right)\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \int\limits_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}} q^{-2k+2}t^2 d_qt$$

$$\begin{aligned}
= {} & \sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \left(\frac{[k]_q+q^{k-1}\alpha}{q^{k-1}\left([n]_q+\beta\right)}\right)^2 \\
& -2\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k q^{-2k+2}q^{k-1}\alpha\frac{[k]_q+q^{k-1}\alpha}{\left([n]_q+\beta\right)^2} \\
& +\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k q^{-2k+2}\frac{q^{2k-2}\alpha^2}{\left([n]_q+\beta\right)^2} \\
& +\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k q^{-2k+2}\left((1+3\alpha)[2]_q+1\right)\frac{[k]_q+q^{k-1}\alpha}{[3]_q\left([n]_q+\beta\right)^2} \\
& -\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k q^{-2k+2}q^kq^{k-1}\alpha\frac{\left((1+3\alpha)[2]_q+1\right)}{[3]_q\left([n]_q+\beta\right)^2} \\
& +\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \frac{q^2\left(1+3\alpha+3\alpha^2\right)}{[3]_q\left([n]_q+\beta\right)^2} \\[2ex]
= {} & L_n^{\alpha,\beta}(e_2;q,x) - \frac{2\alpha}{[n]_q+\beta}L_n^{\alpha,\beta}(e_1;q,x) + \frac{\alpha^2}{\left([n]_q+\beta\right)^2}L_n^{\alpha,\beta}(e_0;q,x) \\
& +\frac{q\left((1+3\alpha)[2]_q+1\right)}{[3]_q\left([n]_q+\beta\right)}L_n^{\alpha,\beta}(e_1;q,x) - q\alpha\frac{\left((1+3\alpha)[2]_q+1\right)}{[3]_q\left([n]_q+\beta\right)^2}L_n^{\alpha,\beta}(e_0;q,x) \\
& +\frac{q^2\left(1+3\alpha+3\alpha^2\right)}{[3]_q\left([n]_q+\beta\right)^2}L_n^{(\alpha,\beta)}(e_0;q,x) \\
= {} & \frac{[n]_q[m(n)]_q}{q\left([n]_q+\beta\right)^2}\left(1+\alpha_{1,m(n)}(x)\right)\left(1+\alpha_{2,n}(x)\right)x^2 \\
& +\frac{[n]_q\left[[3]_q+q\left((1+3\alpha)[2]_q+1\right)\right]}{[3]_q\left([n]_q+\beta\right)^2}\left(1+\alpha_{1,n}(x)\right)x + \frac{q^2\left(1+3\alpha+3\alpha^2\right)}{[3]_q\left([n]_q+\beta\right)^2}.
\end{aligned}$$

This completes the proof of Lemma 6. ■

Using above Lemma, we can obtain following theorem.

Theorem 7 *Let* $f \in C[0,b]$, *then*

$$\lim_{n\to\infty} L_n^{*(\alpha,\beta)}(f;q,x) = f(x)$$

uniformly on $[0,b]$.

3 Rate of convergence

$B_{\rho_\gamma}(\mathbb{R}_+)$, the weighted space of real valued functions f defined on $\mathbb{R}_+$ with the property $|f(x)| \leq M_f \rho_\gamma(x)$ where $\rho_\gamma(x) = 1 + x^{\gamma+2}$ and M_f is constant depending on the function f. We also consider the weighted subspace $C_{\rho_\gamma}(\mathbb{R}_+)$ of $B_{\rho_\gamma}(\mathbb{R}_+)$ given by

$$C_{\rho_\gamma}(\mathbb{R}_+) := \left\{ f \in B_{\rho_\gamma}(\mathbb{R}_+) : f \text{ continuous on } \mathbb{R}_+ \right\}.$$

The norm in B_{ρ_γ} is defined as

$$\|f\|_{\rho_\gamma} = \sup_{x\in\mathbb{R}_+} \frac{|f(x)|}{\rho_\gamma(x)}.$$

We can give some estimations of the errors $\left| L_n^{*(\alpha,\beta)}(f;q,x) - f(x) \right|$, $n \in N$, for unbounded functions by using a weighted modulus of smoothness associated to the space $B_{\rho_\gamma}(\mathbb{R}_+)$.

We consider

$$\Omega_{\rho_\gamma}(f;\delta) = \sup_{x\geq 0,\ 0<h\leq\delta} \frac{|f(x+h) - f(x)|}{1 + (x+h)^{2+\gamma}}, \quad \delta > 0, \quad \gamma \geq 0. \tag{13}$$

It is evident that for each $f \in B_{\rho_\gamma}(\mathbb{R}_+)$, $\Omega_{\rho_\gamma}(f;.)$ is well defined and

$$\Omega_{\rho_\gamma}(f;\delta) \leq 2\|f\|_{\rho_\gamma}.$$

The weighted modulus of smoothness $\Omega_{\rho_\gamma}(f;.)$ posseses the following properties.

$$\Omega_{\rho_\gamma}(f;\lambda\delta) \leq (\lambda+1)\Omega_{\rho_\gamma}(f;\delta), \ \ \delta > 0,\ \lambda > 0, \tag{14}$$

$$\Omega_{\rho_\gamma}(f;n\delta) \leq n\Omega_{\rho_\gamma}(f;\delta),\ n \in N,$$

$$\lim_{\delta\to 0^+} \Omega_{\rho_\gamma}(f;\delta) = 0.$$

As it is known, weighted Korovkin type theorems have been proven by Gadjiev(see [9]).

Theorem 8 *Let $q \in (0,1)$ and $\gamma \geq 0$. For all non-decreasing $f \in B_{\rho_\gamma}(\mathbb{R}_+)$ we have*

$$\left| L_n^{*(\alpha,\beta)}(f;q,x) - f(x) \right| \leq \sqrt{L_n^{*(\alpha,\beta)}(\mu_{x,\gamma}^2;x)} \left(1 + \frac{1}{\delta} \sqrt{L_n^{*(\alpha,\beta)}(\psi_x^2;x)} \right) \Omega_{\rho_\gamma}(f;\delta),$$

$x \geq 0$, $\delta > 0$, $n \in N$, where $\mu_{x,\gamma}(t) := 1 + (x + |t-x|)^{2+\gamma}$, $\psi_x(t) := |t-x|$, $t \geq 0$.

Proof. Let $n \in N$ and $f \in B_{\rho_\gamma}(\mathbb{R}_+)$. From (13) and (14), we can write

$$\begin{aligned} |f(t) - f(x)| &\leq \left(1 + (x + |t-x|)^{2+\gamma} \right) \left(1 + \frac{1}{\delta} |t-x| \right) \Omega_{\rho_\gamma}(f;\delta) \\ &= \mu_{x,\gamma}(t) \left(1 + \frac{1}{\delta} \psi_x(t) \right) \Omega_{\rho_\gamma}(f;\delta). \end{aligned}$$

Taking into account the definition of $q-$ integration, we get

$$\int_{q\left(\frac{[k]_q + q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q + q^k\alpha}{[n]_q+\beta}} f\left(q^{-k+1}t\right) d_qt = q^{k-1} \int_{\frac{[k]_q + q^{k-1}\alpha}{q^{k-2}\left([n]_q+\beta\right)}}^{\frac{[k+1]_q + q^k\alpha}{q^{k-1}\left([n]_q+\beta\right)}} f(t)\, d_qt. \tag{15}$$

Consequently, the operators $L_n^{*(\alpha,\beta)}$ can be expressed as follows

$$L_n^{*(\alpha,\beta)}(f;q,x) = \left([n]_q + \beta \right) \sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!} (-x)^k q^{k-1} \int_{\frac{[k]_q + q^{k-1}\alpha}{q^{k-2}\left([n]_q+\beta\right)}}^{\frac{[k+1]_q + q^k\alpha}{q^{k-1}\left([n]_q+\beta\right)}} f(t)\, d_qt.$$

By using the Cauchy-Schwartz inequality and (15), we obtain

$$\begin{aligned} & \left| L_n^{*(\alpha,\beta)}(f;q,x) - f(x) \right| \\ \leq & \left([n]_q + \beta \right) \sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!} (-x)^k q^{k-1} \int_{\frac{[k]_q + q^{k-1}\alpha}{q^{k-2}\left([n]_q+\beta\right)}}^{\frac{[k+1]_q + q^k\alpha}{q^{k-1}\left([n]_q+\beta\right)}} |f(t) - f(x)|\, d_qt \\ \leq & \left(L_n^{*(\alpha,\beta)}(\mu_{x,\gamma};x) + \frac{1}{\delta} L_n^{*(\alpha,\beta)}(\mu_{x,\gamma}\psi_x;x) \right) \Omega_{\rho_\gamma}(f;\delta) \\ \leq & \sqrt{L_n^{*(\alpha,\beta)}(\mu_{x,\gamma}^2;x)} \left(1 + \frac{1}{\delta} \sqrt{L_n^{*(\alpha,\beta)}(\psi_x^2;x)} \right) \Omega_{\rho_\gamma}(f;\delta). \end{aligned}$$

■

Lemma 9 *For $m \in N$ and $q \in (0,1)$ we have*

$$L_n^{*(\alpha,\beta)}(e_m;q,x) \leq A_{m,q}\,(1+x^m)\ \ ,\quad x \in \mathbb{R}_+,\quad n \in N,$$

where $A_{m,q}$ is a positive constant depending only on m, α and q.

Proof. For $k \in N$ and $0 < q < 1$ the following inequality holds true

$$1 \leq [k+1]_q \leq 2\,[k]_q\,. \tag{16}$$

Thus, for $m \in N$, from (1) and (16) we get

$$\begin{aligned}
L_n^{\alpha,\beta}(e_m;q,x) &= \sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \frac{1}{q^{km-m}}\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)^m \\
&= \frac{x\,[n]_q}{[n]_q+\beta}\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_{m(n)}(x)\right)(1+\alpha_{k,n}(x))}{[k]_q!}(-x)^k \frac{1}{q^{k(m-1)}}\left(\frac{[k]_q+q^{k}\alpha}{[n]_q+\beta}\right)^{m-1} \\
&\quad +\frac{\alpha}{[n]_q+\beta}\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \left(\frac{[k]_q+q^{k-1}\alpha}{q^{k-1}\left([n]_q+\beta\right)}\right)^{m-1} \\
&\leq \frac{x\,[n]_q}{[n]_q+\beta}\varphi_{m(n)}(x)\left(1+\alpha_{0,n}(x)\right)\left(\frac{1+\alpha}{[n]_q+\beta}\right)^{m-1} \\
&\quad +\frac{x\,[n]_q}{[n]_q+\beta}\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_{m(n)}(x)\right)(1+\alpha_{k,n}(x))}{[k]_q!}(-x)^k \left(\frac{2\,[k]_q+q^{k}\alpha}{q^k\left([n]_q+\beta\right)}\right)^{m-1} \\
&\quad +\frac{\alpha}{[n]_q+\beta}L_n^{\alpha,\beta}(e_{m-1};q,x) \\
&= \frac{x\,[n]_q}{[n]_q+\beta}\varphi_{m(n)}(x)\left(1+\alpha_{0,n}(x)\right)\left(\frac{1+\alpha}{[n]_q+\beta}\right)^{m-1} \\
&\quad +\frac{x\,[n]_q}{[n]_q+\beta}\left(\frac{2}{q}\right)^{m-1} L_{n+1}^{\alpha,\beta}(e_{m-1};q,x)+\frac{\alpha}{[n]_q+\beta}L_n^{\alpha,\beta}(e_{m-1};q,x) \\
&\leq x+\left(\frac{2}{q}\right)^{m-1}\frac{1}{[n]_q+\beta}\left(x\,[n]_q+\alpha q^{m-1}\right)L_{n+1}^{\alpha,\beta}(e_{m-1};q,x) \\
&\leq 2m\left(\frac{2}{q}\right)^{m-1}\frac{1}{[n]_q+\beta}\left([n]_q\left(1+x^m\right)+\alpha^m q^{\frac{m(m-1)}{2}}\right).
\end{aligned}$$

based on the above inequality and by using the mathematical induction over $m \in N$, we obtain

$$L_n^{\alpha,\beta}(e_m;q,x) \leq B_{m,q}\left(1+x^m\right),$$

$x \in \mathbb{R}_+,\ n \in N$, where

$$B_{m,q} := 2m\left(\frac{2}{q}\right)^{\frac{m(m-1)}{2}}\left(1+\alpha^m q^{\frac{m(m-1)}{2}}\right). \tag{17}$$

On the other hand,

$$\begin{aligned} L_n^{*(\alpha,\beta)}(e_m;q,x) &= \left([n]_q+\beta\right)\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \int\limits_{q\left(\frac{[k]_q+q^{k-1}\alpha}{[n]_q+\beta}\right)}^{\frac{[k+1]_q+q^k\alpha}{[n]_q+\beta}} e_m\left(q^{-k+1}t\right)d_qt \\ &= \frac{[n]_q+\beta}{\left([n]_q+\beta\right)^{m+1}}\sum_{k=0}^{\infty} q^{\frac{k(k-1)}{2}} \frac{D_q^k\left(\varphi_n(x)\right)}{[k]_q!}(-x)^k \frac{q^{-km+m}}{[m+1]_q} \\ &\quad \times\left\{\left([k+1]_q+q^k\alpha\right)^{m+1}-q^{m+1}\left([k]_q+q^{k-1}\alpha\right)^{m+1}\right\}. \end{aligned}$$

Since

$$\begin{aligned} & \left([k+1]_q+q^k\alpha\right)^{m+1}-q^{m+1}\left([k]_q+q^{k-1}\alpha\right)^{m+1} \\ = & \left(\left([k+1]_q+q^k\alpha\right)^m+q\left([k+1]_q+q^k\alpha\right)^{m-1}\left([k]_q+q^{k-1}\alpha\right)+...+q^m\left([k]_q+q^{k-1}\alpha\right)^m\right) \\ \leq & (m+1)\left([k+1]_q+q^k\alpha\right)^m \\ \leq & (m+1)\,2^m\left([k]_q+q^k\alpha\right)^m,\ k\in N, \end{aligned}$$

from condition (vi), we can write

$$\begin{aligned} L_n^{*(\alpha,\beta)}(e_m;q,x) &\leq \frac{\varphi_n(x)(m+1)(1+\alpha)^m q^m}{\left([n]_q+\beta\right)^m[m+1]_q}+\frac{2^m(m+1)}{[m+1]_q}L_n^{\alpha,\beta}(e_m;q,x) \\ &\leq A_{m,q}\left(1+x^m\right), \end{aligned}$$

where $A_{m,q}:=\frac{(m+1)(1+\alpha)^m q^m}{[m+1]_q}+\frac{2^m(m+1)}{[m+1]_q}B_{m,q}$ and $B_{m,q}$ is given by (17). ■

Remark 10 *Since any linear positive operator is monotone, from Lemma 9 we can easily see that* $L_n^{*(\alpha,\beta)}(f;q,.)\in B_{\rho_\gamma}(\mathbb{R}_+)$ *for each* $f\in B_{\rho_\gamma}(\mathbb{R}_+),\ \gamma\in N_0$.

Theorem 11 *Let* $f\in B_{\rho_\gamma}(\mathbb{R}_+)$ *be a non-decreasing function, then*

$$\left\|L_n^{*(\alpha,\beta)}(f;q_n,.)-f\right\|_{\rho_{\gamma+1}}\leq K_{\gamma,q_0}\Omega_{\rho_\gamma}(f;\delta_n),$$

where $\delta_n:=\sqrt{\frac{[n]_{q_n}\eta_n+1}{q_n\left([n]_{q_n}+\beta\right)}}$ *and* K_{γ,q_0} *is a positive constant independent on* f *and* n.

Proof. The identities (3)-(5) imply

$$\begin{aligned}
L_n^{*(\alpha,\beta)}(\psi_x^2;q_n,x) &= L_n^{*(\alpha,\beta)}\left((t-x)^2;q_n,x\right)\\
&= \frac{[n]_{q_n}[m(n)]_{q_n}}{q_n\left([n]_{q_n}+\beta\right)^2}\left(1+\alpha_{1,m(n)}(x)\right)\left(1+\alpha_{2,n}(x)\right)x^2\\
&\quad+\frac{[n]_{q_n}\left[[3]_{q_n}+q_n\left((1+3\alpha)[2]_{q_n}+1\right)\right]}{[3]_{q_n}\left([n]_{q_n}+\beta\right)^2}\left(1+\alpha_{1,n}(x)\right)x\\
&\quad+\frac{q_n^2(1+3\alpha+3\alpha^2)}{[3]_{q_n}\left([n]_{q_n}+\beta\right)^2}-2x\left\{\frac{[n]_{q_n}}{[n]_{q_n}+\beta}x\left(1+\alpha_{1,n}(x)\right)+\frac{q_n(1+2\alpha)}{[2]_{q_n}\left([n]_{q_n}+\beta\right)}\right\}+x^2\\
&\leq \frac{\left([n]_{q_n}\eta_n(x)+1+\beta\right)x^2}{q_n\left([n]_{q_n}+\beta\right)}+\frac{2(3\alpha+3)}{q_n\left([n]_{q_n}+\beta\right)}x+\frac{1+3\alpha+3\alpha^2}{q_n\left([n]_{q_n}+\beta\right)}\\
&\leq \frac{9(1+\beta)^2\rho_0(x)}{q_n\left([n]_{q_n}+\beta\right)}\left\{[n]_{q_n}\eta_n(x)+1\right\}
\end{aligned}$$

where $\eta_n(x):=\max\left\{\alpha_{1,n}(x),\alpha_{1,m(n)}(x),\alpha_{2,n}(x)\right\}$.
Since $\eta_n(x)$ converges uniformly to zero, we have $\eta_n=\sup\eta_n(x)$ such that η_n converges to zero as $n\to\infty$. Let $\gamma\in N_0$ and $f\in B_{\rho_\gamma}(\mathbb{R}_+)$ be a fixed function. From Theorem 8 and above inequality, we can write

$$\begin{aligned}
&\frac{\left|L_n^{*(\alpha,\beta)}(f;q_n,x)-f(x)\right|}{\rho_{\gamma+1}(x)}\\
&\leq \sqrt{\frac{L_n^{*(\alpha,\beta)}(\mu_{x,\gamma}^2;q,x)}{\rho_{\gamma+1}^2(x)}}\left(1+\frac{1}{\delta_n}\sqrt{L_n^{*(\alpha,\beta)}(\psi_x^2;q_n,x)}\right)\Omega_{\rho_\gamma}(f;\delta_n)\\
&\leq \sqrt{\frac{L_n^{*(\alpha,\beta)}(\mu_{x,\gamma}^2;q_n,x)\rho_0(x)}{\rho_{\gamma+1}^2(x)}}\left(1+\frac{1}{\delta_n}\sqrt{\frac{9(1+\beta)^2\rho_0(x)}{q_n\left([n]_{q_n}+\beta\right)}\left\{[n]_{q_n}\eta_n(x)+1\right\}}\right)\Omega_{\rho_\gamma}(f;\delta_n)\\
&\leq 12(1+\beta)\sqrt{\frac{L_n^{*(\alpha,\beta)}(\mu_{x,\gamma}^2;q_n,x)}{\rho_{2(\gamma+1)}(x)}}\left(1+\frac{1}{\delta_n}\sqrt{\frac{[n]_{q_n}\eta_n(x)+1}{q_n\left([n]_{q_n}+\beta\right)}}\right)\Omega_{\rho_\gamma}(f;\delta_n).
\end{aligned}$$

Since

$$\begin{aligned}
\mu_{x,\gamma}^2(t) &= \left(1+(x+|t-x|)^{2+\gamma}\right)^2\leq 2\left(1+(2x+t)^{4+2\gamma}\right)\\
&\leq 2\left(1+2^{4+2\gamma}\left((2x)^{4+2\gamma}+t^{4+2\gamma}\right)\right),
\end{aligned}$$

from Lemma 9, we get

$$L_n^{*(\alpha,\beta)}(\mu_{x,\gamma}^2;q,x)\leq\lambda_{\gamma,q_n}^2\rho_{2(\gamma+1)}(x),$$

where $\lambda^2_{\gamma,q_n} = 2^{5+2\gamma}\left(2^{4+2\gamma} + A_{4+2\gamma,q_n}\right)$. Choosing $\delta_n := \sqrt{\frac{[n]_{q_n}\eta_n+1}{q_n\left([n]_{q_n}+\beta\right)}}$ and $K_{\gamma,q_0} := 24(1+\beta)\lambda_{\gamma,q_0}$, where $q_0 := \min\limits_{n\in N} q_n$, the proof is finished. ■

References

[1] U. Abel, V. Gupta, An estimate of the rate of convergence of a Bezier variant of the Baskakov-Kantorovich operators for bounded variation functions, Demonstratio math.,36, 123-136, 2003.

[2] A. Aral, V. Gupta, On the Durrmeyer type modification of the $q-$Baskakov type operators, Nonlinear Analysis 72, 1171-1180, 2010.

[3] Ç. Atakut, On Kantorovich-Baskakov-Stancu type operators, Bull. Cal. Math. Soc., 91(2), 149-156, 1999.

[4] V.A. Baskakov, An example of a sequence of linear positive operators in the space of continuous, DAN, 113, 249-251, 1957.

[5] İ.Büyükyazıcı, Ç. Atakut, On Stancu type generalization of $q-$Baskakov operators,52(5-6), 752-759, 2010.

[6] İ. Büyükyazıcı, Direct and inverse results for generalized $q-$Bernstein polynomials, Far East J. Appl. Math., 34(2), 191 – 204, 2009.

[7] R.A. DeVore, G.G. Lorentz, Constructive Approximation, Springer, Berlin, 1993.

[8] A.D. Gadjiev, Ç. Atakut, On approximation of unbounded functions by the generalized Baskakov operators. Trans. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., 23(1), Math. Mech., 33–42, 2003.

[9] A.D. Gadjiev, Theorems of the type of P. P. Korovkin's theorems, Math. Zametki, 20(5), 781-786, 1976, .

[10] N. K. Govil, V. Gupta, Convergence of $q-$Meyer-König-Zeller-Durrmeyer operators. Adv. Stud. Contemp. Math. (Kyungshang), 19(1), 97–108, 2009.

[11] V. Gupta, Z. Finta, On certain $q-$Durrmeyer type operators, Appl. Math. Comput., 209(2), 415-420, 2009.

[12] V. Gupta, C. Radu, Statistical approximation properties of $q-$Baskakov-Kantorovich operators, Cent. Eur. J. Math., 7(4), 809-818, 2009.

[13] S.C. Jing, H.Y. Fan, $q-$ Taylor's formula with its $q-$ remainder, Comm. Theoret. Phys., 23(1), 117-120, 1995.

[14] V.G. Kac, P. Cheung, Quantum Calculus, Universitext, Springer-Verlag, New York, 2002.

[15] G. M. Phillips, Bernstein polynomials based on the $q-$integers, Ann. Numer. Math., 4, 511-518, 1997.

[16] G. M. Phillips, A survey of results on the $q-$ Bernstein polynomials, IMA Journal of Numerical Analysis, 30, 277–288, 2010.

[17] G.M. Phillips, Interpolation and Approximation by Polynomials, Springer-Verlag, New York, 2003.

[18] C. Radu, Statistical approximation properties of Kantorovich operators based on $q-$integers, Creat. Math. Inform., 17, 75-84, 2008.